Satellite Data Delivery in the IOOS Era

Dave Foley

CoastWatch Coordinator
Joint Institute for Marine and Atmospheric Research
University of Hawaii
And

NOAA Southwest Fisheries Science Center, Environmental Research Division.

> NASA OCRT Meeting Seattle, WA April 11, 2007




Overview

- ▲ Background on IOOS and Remote Sensing
 - Spatial Scales to be a big deal
- ▲ Data Stewardship at National Level
- Data Management at Regional Level
- ▲ Data Delivery Mechanisms
- Illustration by Application
 - The need for basic (L2) data
 - The need for highly derived properties
 - The need for long-term continuity





Background: IOOS and Remote Sensing

- ▲ Initial Workshop in Durham, New Hampshire during October 2006.
- ▲ Generally assumed to be a contribution of the "National backbone". Very few elements of the "backbone" are as yet identified
- ▲ Many requirements request support possible only by airborne sensors.




Higher spatial resolution critical to monitor complex coastal waters





Non-Chesapeake Bay Image





Data Stewardship at the National Level

- ▲ Maintenance of archives
 - CLASS (Boulder, CO and Asheville, NC)
 - "Nodes"
- Reprocessing of climate data records
 - Scientific Data Stewardship Committee
- ▲ Development of New Products
 - User driven (e.g., IOOS)
 - "Hey let's try this"



Regional Development Model

- ▲ Form 'centers of excellence" comprised of regional experts
- ▲ Target specific applications
 - Regional algorithms
 - Integrated products
- Develop and deliver experimental products
- ▲ Transfer technology to National level for possible implementation elsewhere



Data Availability

- ▲ Standard transport of large files via ftp or similar means
- Live Access Server browse capability and data delivery of sub-samples in variety of formats
- Interoperable (machine to machine)
 - OPenDAP
 - Web Services (WSC, WMS, WFS)
- Aggregators
 - THREDDS (inherently supports opendap,wcs)
- ▲ Transition from web-based to client-side data transfers





Client-side Services


- ▲ Client accesses the required data from within their favorite application
 - Matlab
 - IDL
 - ArcGIS
 - Excel (??)
 - R/S+
- All relevant data available by a common interface
- ▲ Allows them to integrate their data without sending it in to some [non-existent] grand data base.



Heceta Bank HAB: A Juan de Fuca eddy analog?








Olive Ridley Turtles


- ▲ Yonat Swimmer, Lianne McNaughton and others (NMFS/PIFSC).
- ▲ Anders Nielson and John Sibert (U. Hawaii/ PFRP).
- ▲ Mike Laurs (NMFS/SWFS



Examining Habitat Preferences




Potential Habitat Maps





Need for Long-term Continuity



Ongoing Efforts

- ▲ Learn to handle L1A and L2 swath data with new transport tools.
- ▲ Form partnerships with end users to develop appropriate products.
- ▲ Develop systematic metadata standards, and the ability to translate between those and various existing standards
- ▲ Develop access tools for common clients



Concerns for the Future

▲ Continuity

- VIIRS (??) - who'll fix it if it flys?

▲ Access

 Would like the DAACS to adopt services such as THREDDS

▲ Archives

- CLASS has a LONG was to go, especially with regards to science-quality data records.
- Will the DAACs be there in the interim, under the new NASA model.

Advocacy

- We continue to build advocacy within the Marine resources community; success stories will be shared with NASA as they emerge.



Contact Information

Dave Foley
NOAA SWFSC - ERD
1352 Lighthouse Ave
Pacific Grove, CA 93950

831.648.0632

dave.foley@noaa.gov

http://coastwatch.pfel.noaa.gov/

